



Question Paper



## MANIPAL ACADEMY OF HIGHER EDUCATION

B.Tech First Semester Sessional Examination September 2024

**ENGINEERING MATHEMATICS - I [MAT 1171]**

**Marks: 30**

**Duration: 90 mins.**

**MCQ**

**Answer all the questions.**

Section Duration: 20 mins

- 1) Using Jacobi's Method with initial approximation  $x^{(0)} = y^{(0)} = z^{(0)} = 0$ ,  
the value of  $x$  in the second iteration of the following system of equations is \_\_\_\_\_

$$\begin{aligned} 2x + y + z &= 5 \\ 3x + 5y + 2z &= 15 \\ 2x + y + 4z &= 8 \end{aligned} \quad (0.5)$$

[0](#) [1.5625](#) [2.5](#) [1.5](#)

- 2) The rank of the matrix  $A = \begin{bmatrix} 3 & 1 & -1 & -5 \\ 0 & 0 & 0 & -2 \\ 0 & 7 & -8 & 1 \end{bmatrix}$  is \_\_\_\_\_ (0.5)

[3](#) [1](#) [2](#) [4](#)

- 3) If two of the eigenvalues of  $A = \begin{bmatrix} 4 & 6 & 10 \\ 3 & 10 & 13 \\ -2 & -6 & -8 \end{bmatrix}$  are 0 and 2, then the eigenvalues of  $A^3$  are \_\_\_\_\_ (0.5)

[0, 2, 4](#) [0, 8, 64](#) [0, 4, 16](#) [0, -2, -4](#)

- 4) Which of the following set of vectors is linearly independent in  $\mathbb{R}^2$  ? (0.5)

[{\(0, -1\), \(2, 0\), \(2, -1\)}](#) [{\(-2, 3\), \(2, -3\)}](#) [{\(6, 0\), \(0, -3\)}](#) [{\(5, 0\), \(-7, 0\)}](#)

5)

If the eigenvector of  $A = \begin{bmatrix} 0 & 5 & -10 \\ 0 & 22 & 16 \\ 0 & -9 & -2 \end{bmatrix}$  is  $X = \begin{bmatrix} -5 \\ -4 \\ 3 \end{bmatrix}$ , then the corresponding eigenvalue of A is \_\_\_\_

4 5 10 19

6) The system of equation  $AX = B$  with n unknowns has unique solution if and only if \_\_\_\_

$rank(A) = rank(A|B) = n$   $rank(A) = rank(A|B)$   $rank(A) \neq rank(A|B)$   $rank(A) = rank(A|B) < n$

7) The value of k such that the vector  $\begin{bmatrix} 5 \\ -2 \\ 4 \end{bmatrix}$  is an eigenvector of the matrix  $A = \begin{bmatrix} 5 & -10 & -5 \\ 2 & k & 2 \\ -4 & -8 & 6 \end{bmatrix}$  is \_\_\_\_

1 14 12 17

8) The integrating factor of the differential equation  $\frac{dy}{dx} - 3y \cot x = \sin 2x$ , to reduce it to an exact differential equation is \_\_\_\_

$\sin x$   $\operatorname{cosec}^3 x$   $\sin^3 x$   $\operatorname{cosec} x$

9) If  $A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 0 \end{bmatrix}$ , then the characteristic equation of A is \_\_\_\_

$\lambda^3 - 3\lambda^2 + 2\lambda = 0$   $\lambda^3 + 3\lambda^2 + 2\lambda = 0$   $\lambda^3 + 3\lambda^2 - 2\lambda = 0$   $\lambda^3 - 3\lambda^2 - 2\lambda = 0$

10) The rank of the matrix  $A = \begin{bmatrix} 2 & 1 & 3 \\ 3 & k & -1 \\ 5 & -4 & 2 \end{bmatrix}$  is 2 when k is \_\_\_\_

-5 4 -3 5

### DESCRIPTIVE

Answer all the questions.

11) Test whether the set of vectors  $B = \{ (2, 2, 1), (1, 3, 1), (1, 2, 2) \}$  forms a basis for  $\mathbb{R}^3$  or not. If so, express the vector  $(3, 1, 1)$  in terms of basis vectors.

12) Solve the differential equation  $(4xy + 3y^2 - x)dx + (x^2 + 2xy)dy = 0$  (3)

13) Find all the eigenvalues and the corresponding eigenvectors of the matrix  $A = \begin{bmatrix} 5 & 4 \\ 1 & 2 \end{bmatrix}$  (3)

14) Find the inverse of the matrix  $A = \begin{bmatrix} 1 & 0 & 2 \\ 2 & -1 & 3 \\ 4 & 1 & 8 \end{bmatrix}$ , using Gauss-Jordan method. (3)

15) Test for consistency, the following system of equations and hence solve by Gauss Elimination method

$$x + y + z = 3$$

$$2x - y - z = 3 \quad (3)$$

$$x - y + z = 9$$

16) Solve the differential equation:  $\frac{dy}{dx} - y \tan x = \frac{\sin x \cos^2 x}{y^2}$  (3)

17) Using Gauss Seidel method with initial approximation  $x_1^{(0)} = x_2^{(0)} = x_3^{(0)} = 0$ , solve the following system of equations

$$-3x_1 + 22x_2 + 2x_3 = 47;$$

$$45x_1 + 2x_2 + 3x_3 = 58 ; \quad (2)$$

$$5x_1 + x_2 + 20x_3 = 67.$$

Carry out 2 iterations up to 3 decimal place accuracy.

18) Solve the differential equation  $(3x^2 \tan y - \cos x)dx + x^3 \sec^2 y dy = 0$  (2)

19) Using Rayleigh power method, find the numerically largest eigenvalue and the

corresponding eigenvector of  $A = \begin{bmatrix} 6 & -2 & 2 \\ -2 & 3 & -1 \\ 2 & -1 & 3 \end{bmatrix}$  using the initial vector  $X^{(0)} = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$ . (2)

Carry out 2 iterations up to three decimal place accuracy.