

Type: MCQ

Q1. The stationary point of $f(x, y) = x^3 + y^3 - 3x - 12y + 20 = 0$ is (0.5) **CO2 BL-3**

- A. (1,1)
- B. (2,1)
- C. (-1,1)
- D. ****(-1,2)**

Q2. The coefficient of y^2 in the Maclaurin's series expansion of $f(x, y) = e^x \log(1 + y)$ is (0.5) **CO2 BL-3**

- A. **** $-\frac{1}{2}$**
- B. -1
- C. 1
- D. $\frac{1}{2}$

Q3. The circumference of a circle is measured as 28 cm with an error of 0.01 cm. The percentage error in the area is (0.5) **CO1 BL-3**

- A. **** $\frac{1}{14}$**
- B. $\frac{1}{7}$
- C. $\frac{1}{28}$
- D. $-\frac{1}{28}$

Q4. The value $\lim_{x \rightarrow 0} \frac{3e^x - 2e^{2x} - e^{3x}}{e^x + e^{2x} - 2e^{3x}}$ is (0.5) **CO2 BL-3**

- A. **** $\frac{4}{3}$**
- B. $\frac{1}{4}$
- C. $\frac{1}{2}$
- D. $\frac{3}{4}$

Q5. By changing the integral $\int_0^\infty \int_0^\infty e^{-(x^2+y^2)} dx dy$ into polar coordinates, we get (0.5) **CO3 BL-3**

- A. $\int_0^\infty \int_0^\pi e^{-r^2} d\theta dr$
- B. **** $\int_0^\infty \int_0^\pi e^{-r^2} r d\theta dr$**

C. $\int_0^{\infty} \int_0^{\frac{\pi}{2}} e^{-r^2} 2rd\theta dr$

D. $\int_0^{\infty} \int_0^{\frac{\pi}{2}} e^{-r^2} r^2 d\theta dr$

Q6. The value of $\int_0^1 \int_0^x e^{\frac{y}{x}} dy dx$ is (0.5) **CO3 BL-3**

A. $\frac{e-1}{2}$

B. $e - 1$

C. e^{-2}

D. e

Q7. The center and radius of the sphere $x^2 + y^2 + z^2 - 2y - 4z = 11$ are (0.5)

CO1 BL-3

A. Center $(0, -1, -2)$ and $r = 4$

B. ****Center $(0, 1, 2)$ and $r = 4$**

C. Center $(0, 1, -2)$ and $r = 4$

D. Center $(0, -1, 2)$ and $r = 4$

Q8. If the functions $f(x) = x^3$ and $g(x) = x^2$ satisfy all the conditions of Cauchy's mean value theorem in $[1, 2]$ then the value of c is (0.5) **CO2 BL-3**

A. $\frac{5}{3}$

B. **** $\frac{14}{9}$**

C. 1

D. 2

Q9. If $x = u + uv$ and $y = uv$ then the Jacobian $J = \frac{\partial(x,y)}{\partial(u,v)}$ is (0.5) **CO2 BL-3**

A. **** u**

B. $u + 2uv$

C. uv

D. v

Q10. The Maclaurin's series expansion of $y = \sin x$ is (0.5) **CO2 BL-3**

A. $x + \frac{x^2}{2!} + \dots$

B. **** $x - \frac{x^3}{3!} + \dots$**


C. $1 + x + \frac{x^2}{3!} + \dots$

D. $1 - x + \frac{x^3}{3!} + \dots$

Type: DES

Q11. Using triple integration, find the volume of the paraboloid $x^2 + y^2 = 3z$ cut off by the plane $z = 3$. (4) CO3 BL-3

Ans:- Volume = $4 \iiint dx dy dz$ 1/2 M
 $= 4 \int_0^3 \int_0^{\sqrt{9-y^2}} \int_{z=x^2+y^2}^3 dz dy dx$ 1 M
 $= 4 \int_0^3 \int_0^{\sqrt{9-y^2}} \left(\frac{x^2+y^2}{3}\right) dy dx$



By changing to polar coordinates

$= 4 \int_0^{\pi/2} \int_0^3 \frac{r^2}{3} r dr d\theta$ 1 M
 $= 4 \int_0^{\pi/2} \left(\frac{r^4}{12}\right)_0^3 d\theta = 27 \int_0^{\pi/2} d\theta = \frac{27\pi}{2}$ cubic units 1/2 M

Q12. Using Lagrange's method of undetermined multipliers, find the maximum and minimum distances of the point (1, 2, 3) from the sphere $x^2 + y^2 + z^2 = 56$. (4) CO2

BL-3

Ans:- Let $P(x, y, z)$ be any point on the sphere and $A(1, 2, 3)$. Then $AP^2 = (x-1)^2 + (y-2)^2 + (z-3)^2 = f(x, y, z)$
 $\phi: x^2 + y^2 + z^2 - 56 = 0$ 1 M

Lagrange funct, $L = f + \lambda \phi$ 1/2 M

$\frac{\partial L}{\partial x} = 0 \Rightarrow x-1 = \lambda x$ $\frac{\partial L}{\partial y} = 0 \Rightarrow y-2 = -\lambda y$ 1 M

$\frac{\partial L}{\partial z} = 0 \Rightarrow z-3 = -\lambda z \Rightarrow x = \frac{1}{1+\lambda}, y = \frac{2}{1+\lambda}, z = \frac{3}{1+\lambda}$

$x^2 + y^2 + z^2 = 56 \Rightarrow (1+\lambda)^2 = 1/4 \Rightarrow 1+\lambda = \pm 1/2$ 1/2 M

\therefore Stationary points are $P_1(2, 4, 6), P_2(-2, -4, -6)$ 1/2 M

$$\Rightarrow AP_1 = \sqrt{14} = 3.7 \text{ (Min)}, AP_2 = \sqrt{126} = 11.22 \text{ (Max)} \quad \text{--- } \frac{1}{2} \text{ m}$$

II Semester B.Tech (Common to all Branches) Mid Term Examination March 2024

Subject: Engineering Mathematics II (MAT 1271) Date: 18 March 2024 Time: 2:45 pm to 4:45 pm

Q13. Expand $f(x, y) = \sin(xy)$ in powers of $(x-1)$ and $(y-\frac{\pi}{2})$ up to second degree terms. (3) CO2 BL-3

Ans:

$$f(x, y) = \sin(xy) \Rightarrow f(1, \pi/2) = 1$$

$$f_x = y \cos(xy) \Rightarrow f_x(1, \pi/2) = 0$$

$$f_y = x \cos(xy) \Rightarrow f_y(1, \pi/2) = 0$$

$$f_{xx} = -y^2 \sin(xy) \Rightarrow f_{xx}(1, \pi/2) = -\pi^2/4$$

$$f_{yy} = -x^2 \sin(xy) \Rightarrow f_{yy}(1, \pi/2) = -1$$

$$f_{xy} = -xy \sin(xy) + \cos(xy) \Rightarrow f_{xy}(1, \pi/2) = -\pi/2$$

} 2m

\therefore Req'd expansion is,

$$\sin(xy) = 1 - \frac{\pi^2}{8} (x-1)^2 - \frac{\pi}{2} (x-1)(y-\pi/2) - \frac{1}{2} (y-\pi/2)^2 + \dots$$

--- 1m

Q14. Find the equation of the sphere which passes through the circle

$$S: x^2 + y^2 + z^2 - 2x - 3y + 4z + 8 = 0 \quad ; \quad U: x - 2y + z - 8 = 0$$

and has its center on the plane $4x - 5y - z - 3 = 0$. (3) CO1 BL-3

Ans: Eqⁿ of the req'd sphere: $S + kU = 0$

$$\Rightarrow x^2 + y^2 + z^2 + x(-2+k) + y(-3-2k) + z(4+k) + 8 - 8k = 0$$

--- 1 1/2 m

Centre of \otimes is, $(\frac{2-k}{2}, \frac{3+2k}{2}, \frac{-k-k}{2})$ --- 1/2 m

This centre lies on $4x - 5y - z - 3 = 0$

$$\Rightarrow k = \frac{-9}{13} \quad \text{--- } \frac{1}{2} \text{ m}$$


Req'd eqⁿ: $13(x^2 + y^2 + z^2) - 35x - 21y + 43z + 32 = 0$

II Semester B.Tech (Common to all Branches) Mid Term Examination March 2024

Subject: Engineering Mathematics II (MAT 1271) Date: 18 March 2024 Time: 2:45 pm to 4:45 pm

(1/2 M)

Q15. The region of evaluation of the integral $I = \int_0^{1/2} \int_x^{1-x} xy \, dy \, dx$ is shaded in the following diagram.



Change the order of integration and hence evaluate it. (3) CO3 BL-3

Ans:.

$$I = \int_0^{1/2} \int_0^y xy \, dx \, dy + \int_{1/2}^1 \int_0^{1-y} xy \, dx \, dy \quad \text{--- 1M}$$

$$= \frac{1}{2} \int_0^{1/2} y^3 \, dy + \frac{1}{2} \int_{1/2}^1 (1-y)^2 y \, dy \quad \text{--- 1M}$$

$$= \frac{1}{2} \left(\frac{y^4}{4} \right)_0^{1/2} + \frac{1}{2} \left(\frac{y^2}{2} + \frac{y^4}{4} - \frac{2y^3}{3} \right)_{1/2}^1 \quad \text{--- 1/2 M}$$

$$= \frac{1}{48} \quad \text{--- 1/2 M}$$

Q16. Evaluate $\int_0^a \int_y^a y\sqrt{x^2+y^2} dx dy$ where $a > 0$ by changing to polar coordinates. (3)

C03 BL-3

Ans: put $x = r\cos\theta$, $y = r\sin\theta$ } $\frac{1}{2}M$
 $dx dy = r dr d\theta$


Let $I = \int_0^a \int_y^a y\sqrt{x^2+y^2} dx dy$

$= \int_{\theta=0}^{\pi/4} \int_{r=0}^{a\sec\theta} r\sin\theta \cdot r \cdot r dr d\theta$ — $1M$


$= \int_{\theta=0}^{\pi/4} \sin\theta \left(\frac{r^4}{4}\right)_0^{a\sec\theta} d\theta$ — $\frac{1}{2}M$

$= \frac{a^4}{4} \int_{\theta=0}^{\pi/4} \sin\theta \sec^4\theta d\theta$ } $\frac{1}{2}M$

$= \frac{a^4}{4} \times \frac{(2\sqrt{2}-1)}{3} = \frac{a^4(2\sqrt{2}-1)}{12}$



Q17. Using double integrals, find the area of the shaded region in the given diagram, bounded by the curves $r = 4(1 - \cos \theta)$ and $r = 4 \sin \theta$.



(3) CO3 BL-3

Ans: Area = $\int_{\theta=0}^{\pi/2} \int_{r=0}^{r=4(1-\cos\theta)} r dr d\theta$ + $\int_{\theta=\pi/2}^{\pi} \int_{r=0}^{r=4\sin\theta} r dr d\theta$ — (1M)

$= \int_{\theta=0}^{\pi/2} 8(1-\cos\theta)^2 d\theta + \int_{\theta=\pi/2}^{\pi} 8\sin^2\theta d\theta$ — (1M)

$= 8 \int_{\theta=0}^{\pi/2} (1 - 2\cos\theta + \cos^2\theta) d\theta + 8 \int_{\theta=\pi/2}^{\pi} \frac{(1 - \cos 2\theta)}{2} d\theta$

$= 8 \left(\theta - 2\sin\theta \right) \Big|_0^{\pi/2} + 8 \cdot \frac{1}{2} \times \frac{\pi}{2} + 4 \left(\theta - \frac{\sin 2\theta}{2} \right) \Big|_{\pi/2}^{\pi}$ — (1/2 M)

$= 8 \left(\frac{\pi}{2} - 2 \right) + 2\pi + 4 \left(\frac{\pi}{2} \right)$

$= \underline{\underline{8\pi - 16 \text{ sq. units}}}$ — (1/2 M)

